

Before

After

Unretouched clinical photos taken at maximum frown before treatment with BOTOX® Cosmetic and taken at maximum frown after treatment with BOTOX® Cosmetic at day 14. Your own results may vary.

- *Results may vary. In clinical trials, at day 7, 74% of patients demonstrated "None" or "Mild" glabellar line severity at maximum frown. At day 30, 80% of patients demonstrated the same.*

Everyday people. Everyday expressions.

BOTOX® Cosmetic is approved for the temporary treatment of moderate to severe frown lines between the brows in people ages 18 to 65.

Important Safety Information: *Serious heart problems and serious allergic reactions have been reported rarely. If you think you're having an allergic reaction or other unusual symptoms, such as difficulty swallowing, speaking, or breathing, call your doctor immediately. The most common side effects following injection include temporary eyelid droop and nausea. Localized pain, infection, inflammation, tenderness, swelling, redness, and/or bleeding/bruising may be associated with the injection. Patients with certain neuromuscular disorders such as ALS, myasthenia gravis, or Lambert-Eaton syndrome may be at increased risk of serious side effects.*

Express yourself today.

Ask your doctor if BOTOX® Cosmetic is right for you.

BOTOX®
—Cosmetic
Botulinum Toxin Type A

Full prescribing information has been provided to your doctor.

 **ALLERGAN**
THE SCIENCE OF REJUVENATION™

©2007 Allergan, Inc., Irvine, CA 92612 ® and ™ marks owned by Allergan, Inc. www.BotoxCosmetic.com 1-800-BOTOXMD Re-order: APC011582007 701525